

Pathogens are right within reach with a standard glove box

Only 42%

of staff performed hand hygiene before gloving yet, 100% believed they were compliant.¹

75% of gloves


sampled from patient rooms were contaminated with at least 1 species of bacteria.²



Use these clean glove-pull criteria to reduce HAIs



Practice proper hand hygiene as defined by the APIC



Prevent unused gloves from exposure to airborne bacteria and viruses



Retrieve gloves with the least amount of physical contact possible

Medline's SmartBoX™ reduces contamination by 44%.³

SmartBoX supports clean glove-pull criteria—pushing gloves to the box opening.



Scan to learn more



Medline Industries, LP | Three Lakes Drive, Northfield, IL 60093 | 1-800-MEDLINE (633-5463)

REFERENCES: 1. Baloh, J., Thom, K. A., Perencevich, E., Rock, C., Robinson, G., Ward, M., Herwaldt, L., & Reisinger, H. S. (2019). Hand hygiene before donning nonsterile gloves: Healthcare workers' beliefs and practices. American journal of infection control, 47(5), 492–497. <https://doi.org/10.1016/j.ajic.2018.11.015> 2. Diaz, M. H., Silkaitis, C., Malczynski, M., Noskin, G. A., Warren, J. R., & Zembower, T. (2008). Contamination of Examination Gloves in patient Rooms and Implications for Transmission of Antimicrobial-Resistant Microorganisms. Infection Control & Hospital Epidemiology, 29(1), 63–65. doi:10.1086/524338 3. Data on file. © 2024 Medline Industries, LP. All rights reserved. Medline and SensiCare are registered trademarks and SmartBoX is a trademark of Medline Industries, LP. / MKT19W6183057 / e24379 / 49